

Ultrasonography-Guided Biopsy Techniques: Current Trends, Challenges, and Future Directions

Authors:

Deepanshu Chauhan¹, Divyansh Singh¹, Md. Imran¹, Abhishek Pandey¹, Priyanshu Singh¹, Anmol Kumar², Pradeep Kumar², Ankush Verma³, Srishti⁴, Amit Pratap Singh Chouhan^{4*}

¹Student, Department of Radiology, School of Allied Health Sciences, Galgotias University.

²Demonstrator, Department of Radiology, School of Allied Health Sciences, Galgotias University.

³Assistant Professor, Department of Radiology, Sharda School of Allied Health Sciences, Sharda University.

⁴Assistant Professor, Department of Radiology, School of Allied Health Sciences, Galgotias University.

Corresponding Author:

Amit Pratap Singh Chouhan, Assistant Professor, Department of Radiology, School of Allied Health Sciences, Galgotias University, Greater Noida, India. Orcid ID - 0000-0002-1407-5033

<https://doi.org/10.5281/zenodo.15272855>

Article Received: 09-March-2025, Revised: 29-March-2025, Accepted: 19-April-2025

ABSTRACT:

Ultrasonography has become a crucial part of the majority of interventional radiology departments in recent years. For biopsy procedures, ultrasonography offers reliable and efficient guidance. By allowing the radiologist to always keep the needle's path inside the ultrasonic beam, needle-guidance systems can make biopsy procedures easier for radiologists with little experience. The needle can be confidently advanced on target in this way. Skilled interventional radiologists can employ the freehand technique, which gives them more freedom in terms of skin puncture and needle path, enabling them to be creative in achieving challenging targets. The ability to see the needle tip at all times during the insertion and sampling process is the primary benefit of ultrasonography over computed tomography in image-guided biopsy. When the lesion is very deep or when the lesion is obscured by highly reflective surfaces (such as bone or a gas-filled colon), ultrasound is not very useful for guiding biopsies in obese patients. Ultrasound guided biopsy is a type of diagnostic procedure in which the professional extracts a piece of tissue from infected area for examination due to which we are able to identify the type of disease, the level of damage or infection. The main outcome after the research paper is that we get information related to biopsy which include procedure, benefits, risk, side-effects and post procedural care and conclusion. The aim of the research is to give an ideal knowledge regarding biopsy and ultrasound use in biopsy procedure with the help of suitable examples. A minimally invasive, real-time, safe and efficient method of diagnosing peripheral lung lesions is ultrasonic guided lung biopsy. Clinically, USLB is now rather common. USLB lacks extensive sample studies, therefore its accuracy is poor; moreover, specimens from USLB are hardly used for microbial cultures. After reading this research paper it will help you to fully grasp the ideas of the technique, including the real-time imaging features that support exact needle placement in desired tissues. Such revelations also relate to the clinical uses of this minimally invasive technique, especially for the diagnosis of several diseases including cancer.

Keywords: Ultrasonography, Interventional Radiology, Biopsy, Medical Imaging, Diagnostic Procedure, Invasive.

INTRODUCTION:

A biopsy is a procedure in which we cut a small piece of tissue from infected area using a device called biopsy gun for microscopic analysis or identification and resolution of the disease. When this procedure is analysed or observed under ultrasound due to which it provides preciseness to the biopsy scan. Using some modalities, we can detect precise abnormal tissue, modalities like ultrasound scans, magnetic resonance imaging (MRI) and computed tomography (CT) scans. The biopsy majorly used with benign and malignant tumours by which it can cause cancer in future. A

crucial diagnostic technique for solid pancreatic masses is image-guided biopsy. The purpose of this study is to assess the procedure's accuracy and potential complications (Davis, M. *et al.* (2002)). An ultrasound-guided biopsy is a technique whereby the radiologist locates the particular site and removes the tissue sample under direction from an ultrasound. Ultrasounds create images of an organ or tissue by means of sound waves. High-frequency sound waves produced by the ultrasonic unit enter the body and reflect back upon contacting some body structures. After that, depending on frequency, loudness, and

return time, a computer generates images from these returning sound waves. A tiny hollow needle is also placed into the same site concurrently to get the tissue sample. The site precision of an ultrasonic-guided biopsy is one of its advantages over a traditional one. Moreover, it is less intrusive since the biopsy employs a small needle, so minimizing organ distortion and generating very little scarring.

History of ultrasound- guided biopsy:

Ultrasound-guided biopsy has really changed the game in medical diagnostics, making tissue sampling safer and more accurate. This technique has a history that goes back several decades and has evolved along with technology. Let's take a closer look at how it all started and developed:

History of inventor:

Somewhere back in the 1700s, an Italian priest and physiologist named Lazzaro Spallanzani started some pretty interesting experiments with ultrasound to figure out how bats navigate in the dark. In a rather shocking move, he blinded these bats with a red-hot needle (can you imagine if that went through today's ethics committees?). He found that even in complete darkness, the bats could dodge obstacles while flying around, but when their mouths were covered, they collided with everything in sight. He concluded that bats rely on their ears way more than their eyes to "see," or at the very least, to judge distances. It wasn't until two Harvard students Donald Griffin and Robert Galambos came along and used a sonic detector to capture the ultrasound sounds that bats were making while they flew that the science behind all this started to come together. We've since learned that other animals, like rodents, dogs, moths, dolphins, whales, and even frogs, use ultrasound for all sorts of reasons, like communicating, dodging predators, and hunting. This ability to send out short bursts of sound called echolocation happens across a range of frequencies, from around 1 kHz in birds to about 2 kHz in whales.

The Early Days: Biopsy Without Imaging:

The idea of a biopsy taking a tissue sample to diagnose conditions dates back to the 19th century. Back then, doctors had to rely on their own skills and experience to guide them, doing procedures without any imaging. This "blind" approach often led to issues, like incorrect samples and accidental damage to nearby tissues. Then came the introduction of imaging, which was a real turning point. In the mid-20th century, X-rays were among the first imaging techniques used, followed by computed tomography (CT). While these tools helped improve accuracy, they had their downsides, like radiation exposure and not being able to offer real-time guidance.

The Rise of Ultrasound in Medicine:

Ultrasound started gaining popularity in the mid-20th century, initially focusing on fields like obstetrics and cardiology. Its non-invasive nature, ability to produce real-time images, and lack of harmful radiation made it an attractive option for other medical uses. By the 1970s and 1980s, doctors began to explore its potential for guiding needle biopsies (AJR Am J Roentgenol. 2007).

A key early application of ultrasound-guided biopsy was in the liver. Given the liver's complicated blood vessel structure, traditional biopsy methods were often tricky and risky. Ultrasound allowed doctors to place needles with precision and considerably cut down on complications such as bleeding or missing tumours.

The Advancement of Ultrasound-Guided Techniques:

The 1980s and 1990s saw major improvements in ultrasound technology, including sharper images and portable machines. These enhancements opened the door for ultrasound-guided biopsy to be used in various organs like the kidneys, thyroid, breast, and lymph nodes.

During this time, two main techniques became popular:

1. **Fine-Needle Aspiration (FNA):** This approach uses a thin needle to collect cells for examination. It's minimally invasive and often used for thyroid, lymph node, and breast biopsies.
2. **Core Needle Biopsy (CNB):** This method employs a larger needle to obtain a tissue sample for more detailed analysis. CNB offers more comprehensive insights, making it a go-to for diagnosing cancers and other complicated conditions.

The choice between FNA and CNB really depends on the specific situation, but both benefit greatly from ultrasound guidance.

Modern Uses and Innovations:

Nowadays, ultrasound-guided biopsy is a fundamental tool in diagnostic medicine. It's used to assess lesions in the liver, kidneys, pancreas, prostate, thyroid, and breast, among other areas. This technique is especially essential in oncology, helping to spot cancer early and accurately.

Recent innovations have taken its capabilities even further:

- **Fusion Imaging:** This combines real-time ultrasound with existing CT or MRI images, allowing for even more precise targeting of lesions that might not be clearly visible on ultrasound alone.
- **Elastography:** This ultrasound technique measures tissue stiffness, giving doctors extra information to help guide biopsy decisions, particularly in liver and breast diseases.

Rationale for a biopsy under ultrasound:

The early imaging only reveals abnormalities within the body therefore it needs more imaging for more investigation about the infected area. The biopsies-guided by ultrasound provide more clarity in form of scan, provides more visible area to search for. Ultrasound guided biopsy results in finding Tumours like benign and malignant. However, the finding or identifying the malignant tumours ensures timely check up to control inhibition and timely referrals to the doctor for tumours management (Choi, A.Y. *et al.* (2022)).

Process of decision making:

The process of decision making is necessary and to be performed during or before ultrasound guided biopsy the decision is collectively made by the diagnostic team and patient, patient family members. The diagnostic team include consultant radiologist, clinical nurse specialist and patient, patient family members. The diagnostic team is made on the basis of patient disease, history and investigations before times. Patient collaboration and consent towards the check up and scan can help in continuing the procedure effortlessly and without any obstacles. Decision-making during an ultrasound-guided biopsy process requires a series of well-coordinated steps to obtain accuracy and avoid harm to the patient. This starts with an assessment of the clinical need for the biopsy through previous imaging or symptoms, and then informing the patient of the advantages, complications, and options of the procedure to get their informed consent (Facciorusso, A. *et al.* (2021)). The expert plans the procedure attentively, including the size, location, and accessibility of the lesion. Real-time ultrasound imaging is used during the biopsy to guide the needle to the target tissue, thereby greatly eliminating the possibility of error. Post-procedure monitoring of the patient for complication and proper post-procedure care are also observed, while the tissue sample is submitted for pathological examination to establish or confirm the diagnosis. This systematic way allows the process to be effective and focused on the health of the patient.

Preparation before the biopsy:

- Patients should dress in comfortable clothing and take off any jewellery or accessories prior to visiting for the procedure.
- They should report to their doctor about any allergy history.
- Patients on blood thinners or Aspirin might be required to discontinue these medications for three to five days prior to the procedure to prevent unwanted bleeding.
- No fasting is required for breast, thyroid, or liver biopsies. Patients will be instructed to

fast for any abdominal biopsy six to eight hours prior to the procedure.

- They may have water up to two hours' prior to the procedure.
- Diabetic patients should also notify the doctor as they may be given special instructions regarding the procedure.
- The patient should notify the doctor of their prescription, supplements, or over-the-counter medications.
- Patients who are about to undergo a breast biopsy are advised against the use of deodorant, talcum powder, and lotions on the breast or armpits.
- Physicians can recommend some blood tests prior to the procedure. Patients should bring their recent reports and prescription medication for the procedure (Fallenberg, E.M. (2022)).

Procedure details:

The biopsy procedure is performed by the department of imaging radiology who are skilled in image guided techniques.

How this procedure performed:

Step:1 the patient lies on an ultrasound table and the procedural site (infected site) is sterilized.

Step:2 local anaesthesia is injected into the procedural area for numbing the area.

Step:3 own needle inserting, the extraction of sample tissue from procedural site under the guidance of ultrasound.

Duration of procedure is generally 30 to 45 minutes.

During the sample extraction, patience may hear a 'clunking' sound caused by biopsy equipment, which is normal in biopsy procedure. After the completion of the biopsy procedure, the procedural site is covered with dressing or a plaster so that no contamination can cause (Bass, E.J. *et al.* (2022)).

Equipment used in ultrasound biopsy procedure:

Ultrasound-guided biopsy procedures use specialized equipment to ensure precision, safety, and effectiveness. Let's take a closer look at the key tools and materials involved:

Ultrasound Machine:

This machine is the heart of the procedure. It generates real-time images of the area being targeted, which helps healthcare providers accurately guide the biopsy needle. Many modern machines come with features like Doppler imaging, electrography, and fusion imaging to improve visualization.

Transducer:

The transducer is a handheld device that sends and receives ultrasound waves. It varies in shape and size depending on what's being examined. For biopsies, high-frequency linear transducers are typically used for surface tissues, while curved transducers work better for deeper areas.

Biopsy Needles:

Biopsy needles team up to extract tissue samples with minimal invasiveness. Here are the main types:

Fine-Needle Aspiration (FNA) Needles: These thin, hollow needles collect cells for cytological analysis.

Core Needle Biopsy (CNB) Needles: Larger needles that grab tissue samples for histological examinations.

Cutting Needles: These specialized needles have serrated or beveled tips for precise tissue cutting.

Needle Guides and Sterile Drapes and Covers.

Fig: Showing the Labelled Diagram of Biopsy Needle.





Fig: Showing the Equipment's which are used during the biopsy.



Needle guides are attachments for the transducer that help align the needle with the ultrasound beam. They ensure the needle is placed accurately and lower the chance of complications. Using sterile drapes or sheaths over the transducer and other equipment helps maintain a clean environment, reducing the risk of infection during the procedure.

Local Anaesthetics:

Local anesthetic are applied so that the patients experience no discomfort during tests such as biopsies. Such drugs interfere by inhibiting temporarily nerve signals at the target zone, so no pain can be sensed.

Such drugs are injected through minute syringes and fine needles so that patients get the minimum inconvenience while injections take place. Such drugs will differ according to what is needed from the patients as well as on the nature of the treatment to be provided. Local anaesthesia is rapid inaction, enabling physicians to conduct biopsies effectively while maintaining patient comfort. Patients remain awake during the procedure but feel no pain, facilitating quicker recovery and minimizing risks from general anaesthesia. Vacuum-assisted devices. In certain situations, vacuum-assisted biopsy devices are handy.

They use suction to draw tissue into the needles, allowing for larger samples with fewer insertions.

Specimen Containers and Monitoring Equipment:

Once the tissue sample is taken, it goes into sterile containers for transport to the pathology lab. These containers are usually filled with a preservative solution to keep the sample intact. For procedures involving deeper tissues or sensitive areas, monitoring equipment like pulse oximeters and blood pressure cuffs might be used to ensure safety.

Imaging accessories:

Imaging accessories, especially fusion imaging systems, are now an essential part of ultrasound-guided biopsies, offering increased accuracy and diagnostic dependability. Fusion imaging combines the real-time ultrasound information with pre-acquired CT or MRI imaging, blending the best in both imaging modalities. Although ultrasound offers real-time imaging and visualization of soft tissues, CT and MRI offer the precise anatomical details and an overall view of the area under evaluation. This combination enables one to comprehend intricate structures more clearly and identify target areas more effectively.

When applied to biopsies, for which accuracy of pinpoint is a prerequisite in order to acquire tissue from known lesions or abnormalities, fusion imaging proves to be a beneficial guide. With superimposition of images of various modalities, clinicians can precisely align the ultrasound probe with the site to be sampled, thus preventing sampling errors and complications. It proves particularly useful in the instance of small lesions, those situated deep, or in close proximity to structures that are vital.

The employment of these new imaging devices not only increases procedural success rates but also increases patient safety by reducing invasive attempts. Fusion imaging systems are a major leap in the technology of interventional radiology, where biopsies are efficient and accurate, resulting in enhanced diagnostic and therapeutic outcomes. Documentation Tools. Healthcare providers often rely on software integrated with the ultrasound machine to document the procedure and store images for later reference (Kaplan I, Oldenburg NE, 2002).

Preparation and consideration:

Patient may require a blood test for evaluation of coagulation parameters. Medication such as antiplatelet or anticoagulants maybe required for the patients on medical advice by the doctors. Patient or encourage any kind of information related to themselves before the procedure, so that no side effect shown after the procedure and patient can go happily.

Post-procedural care:

After the biopsy procedure, patients are monitored or under surveillance in a recovery area 4 up to 2 hours to see in case of post procedural allergy or any kind of infection. Responsible adult or aged patients can go home but by taking all precautions as needed after the procedure. Normal medicine can be resumed unless or otherwise advised by doctors. The site of biopsy can make you feel pain and the pain get lighten or decreased by taking analgesics.

Analgesics are also known as painkillers. They are medications that relieve different types of pain from head to injuries to arthritis. Anti-inflammation analgesics can also reduce inflammation.

The dressing should be tied at the site for 24 hours, after that it can be removed. If there is redness, swelling or any kind of discharge after removing dressing have to consult to the doctor immediately. Patient are advised note to drive till 24 hours due to which the tissue extracted from site can healed (Lubner, M.G. *et al.* (2021)).

Ultrasound-guided biopsy has numerous Benefits and some Risks:

Benefits:

Ultrasound-guided biopsies have a variety of advantages that make them a first choice in diagnostic interventions. One of the key benefits is that they are minimally invasive. Ultrasound-guided biopsies are conducted with a thin needle, which avoids the necessity for surgical cuts. This leads to shorter recovery times, less pain, and fewer risks of complications than with conventional surgery. Patients are usually able to return to their usual activities shortly after the procedure, which makes it more convenient.

Real-time imaging is another important advantage. Ultrasound offers real-time images that permit accurate needle placement, which allows tissue samples to be taken precisely from the site of interest. This reduces sampling errors and increases diagnostic accuracy. Real-time visualization is especially beneficial in sensitive procedures where accuracy is paramount, for example, biopsy of small or deeply seated lesions (JAMA. 2015).

The utility of ultrasound-guided biopsies should also be mentioned. The flexibility of this process is impressive because it can be used on diverse tissues such as the breast, liver, thyroid, and lymph nodes. Being so versatile means that it has applicability for different medical disorders and body areas, making diagnostics simpler across numerous specialties.

In addition, ultrasound-guided procedures also prevent radiation exposure, as opposed to CT-guided biopsies. This factor is particularly valuable for patients who need repeated diagnostic tests or those who are at increased risk of complications due to radiation. The lack of radiation helps in the safety and sustainability of this technique.

Cost-effectiveness makes ultrasound-guided biopsies even more appealing. Ultrasound-guided biopsies tend to be cheaper than methods involving CT or MRI-guided biopsies, making them viable for a larger number of patients. Coupling affordability, safety, and efficiency makes ultrasound-guided biopsies an ideal option in medical diagnostics.

Risks:

Although ultrasound-guided biopsies are safe, they are not without risk. One such risk is bleeding or formation of a hematoma at the biopsy site. This occurs in very few instances and generally is minor, although it may take extra treatment to be resolved. Pressure applied to the biopsy site after the procedure generally prevents this.

Another point to consider is the minor risk of infection. Since the needle penetrates the skin, there is always

some small chance that bacteria may enter and cause an infection. However, the use of sterile techniques and proper aftercare greatly minimizes this risk.

In a few exceptional cases, there can be accidental damage to the surrounding structures. For instance, in some biopsies, the needle might inadvertently hurt surrounding tissues like the chest wall or even lungs in situations where the procedure is very close to these structures. These complications are extremely uncommon given the accurate real-time imaging through ultrasound, allowing for cautious steering.

Although these risks are rare, awareness of them ensures that patients are able to make informed choices when discussing the procedure and safety with their doctors. Even in light of these occasional complications, most patients' benefits of precise diagnosis and minimal treatment outweigh the risk involved.

Ultrasound-guided biopsies exist in a number of different types, which are specialized according to specific diagnosis requirements:




Fig : Showing the Different types of Ultrasound Guided Biopsies Techniques.

Fine-Needle Aspiration (FNA): A thin needle is utilized for sampling cells or fluid from an abnormal location to be studied. It's an extremely small-incision technique frequently utilized in the case of thyroid or lymph node biopsies. **Fine-Needle Aspiration** is a minimally invasive technique employed to aspirate cells or fluid from a suspicious mass, like a lump or abnormal tissue, for diagnostic testing. It is most often done under ultrasound guidance to provide accuracy in targeting the area of concern (Takano, Y. *et al.* (2021)).

Procedure:

Preparation: The patient is typically placed comfortably, and the skin over the site of the biopsy is cleaned and sterilized. Local Anaesthesia can be used to minimize discomfort.

Ultrasound Guidance: A miniature ultrasound probe is employed to see the region of interest in real time. This guides the physician precisely to place the fine needle at the correct position.

Aspiration: A thin, empty needle connected to a syringe is placed within the abnormal tissue or mass. Gentle suction is applied by the physician to trap cells or fluid inside the syringe.

Sample Collection: After a sufficient sample is collected, the needle is withdrawn. The material collected is prepared on slides or in a specimen container for cytological evaluation by a pathologist.

Uses of fine needle aspiration:

- **Disorders of the Thyroid:** FNA is routinely employed to assess thyroid nodules or lumps to determine if they are benign (non-cancerous) or malignant (cancerous).

It aids in the diagnosis of disorders such as thyroid cancer, Hashimoto's thyroiditis, or other thyroid disorders.

- **Evaluation of Breast Lump:** For breast lumps, FNA separates cysts (fluid-filled sacs) from solid masses, possibly detecting malignancies. it is frequently the initial diagnostic tool for women presenting with palpable breast lumps.
- **Lymph Node Analysis:** Increased lymph nodes are usually assessed with FNA to check for infection, lymphoma, or metastatic cancer (cancer spread from other locations in the body). This is especially helpful to direct cancer staging and treatment.
- **Liver and Pancreas:** FNA is applied to sample suspicious liver or pancreatic abnormalities that have been identified on imaging tests (e.g., ultrasound or CT scans). It assists in the diagnosis of primary cancers (e.g., hepatocellular carcinoma) or metastases.
- **Lung and Pleural Disorders:** For lung lesions or pleural abnormalities, FNA diagnoses cancers, infections, or inflammatory diseases. it is commonly performed under the guidance of endoscopic ultrasound to increase accuracy in these areas (Facciorusso, A. *et al.* (2022)).

Core Needle Biopsy: A wider, hollow needle is inserted to extract a tiny cylinder of tissue, offering greater information than FNA. It's most often applied for breast, liver, or kidney biopsies.

Vacuum-Assisted Biopsy:

It applies a vacuum-assisted device that extracts several samples of tissue by a single needle puncture, minimizing the repetition of punctures. Vacuum-assisted biopsy (VAB) is a minimally invasive technique applied to acquire tissue samples for diagnostic purposes, especially in the case of breast abnormalities. This innovative method utilizes a hollow needle attached to a vacuum device, enabling accurate and effective removal of tissue samples. In contrast to conventional biopsy techniques, VAB involves a minimal incision, minimizing pain and scarring.

Throughout the procedure, imaging guidance like ultrasound or mammography is employed to precisely identify the area of interest. The vacuum mechanism allows for the removal of several tissue samples via a single insertion, which makes the process faster and less invasive. This is particularly useful for sampling larger or multiple lesions without the necessity of repeated needle insertions.

VAB is universally utilized in diagnosing benign as well as malignant conditions such as breast lump,

calcification, and other abnormal findings in screening procedures. VAB is sought after because of its ability to obtain good quality tissue specimens and its promise in patient comfort and safety. Outpatient performance of the procedure permits patients to recover from the same in the course of their routine activities in short order.

Apart from its use in diagnostics, VAB can also be employed therapeutically to excise small benign lesions completely. Its accuracy and minimally invasive approach make it a prized asset in contemporary medicine, providing accurate results with minimal disruption to the patient's physical and emotional state.

Endoscopic Ultrasound-Guided Biopsy: Blends endoscopy with ultrasound to reach difficult-to-visualize areas, like the pancreas or gastrointestinal tract, to sample tissue. Endoscopic Ultrasound-Guided Biopsy (EUS-guided biopsy) is a cool little medical trick that's like having a tiny camera and an ultrasound wand on a flexible tube to grab tissue samples from parts of your body that are tough to get to, like your pancreas, liver, gut, and lymph nodes. It's a big help when regular biopsies can't cut it because of where the weird stuff is hanging out (Madhok, I.K., Parsa, N. and Nieto, J.M. (2022)).

Procedure:

They'll usually make sure you're all relaxed with some sedation or full-on sleepy juice (general anesthesia). They've got this thing called an endoscope, which is basically a super-thin tube with a camera attached, and it's got a fancy ultrasound probe on the end. They slide this endoscope through your mouth, down into your digestive system so that the probe can show them really detailed pics of your insides. Then, when they spot the area, they want to check out, they use the ultrasound to guide a teeny-tiny needle through the tube and into the tissue that's acting up. They snatch a few samples and send them off to the lab folks to figure out what's going on, like if it's something bad like cancer or not.

Applications:

Pancreas stuff: Like if you've got some suspicious spots or cysts that are playing hide and seek.

Liver and gallbladder: To peek at and sample any funky tissue that's causing trouble.

Gut issues: If you've got a mass in your stomach or intestines that needs a closer look.

Chest lymph nodes: To check if there's anything fishy with them, like lymphoma or if something else has spread there.

Benefits of Endoscopic Ultrasound-Guided Biopsy (EUS-guided biopsy):

Minimally Invasive: EUS-guided biopsy is different from traditional surgical methods. It doesn't need large

cuts or incisions. Instead, doctors use an endoscope that enters through natural openings in the body, like the mouth. This approach is much gentler on the body. The procedure allows for a quicker recovery and less discomfort afterwards because it's less invasive.

Real-Time Imaging: The use of ultrasound with the endoscope provides clear, real-time images. This allows doctors to place the needle precisely where needed. As a result, the test is more accurate, and there are fewer mistakes in sampling. (Sountoulides, P. *et al.* 2021).

Reach Areas: EUS-guided biopsy is especially useful for reaching areas like the pancreas, liver, lymph nodes, and structures in the chest. These places are often hard to reach with other biopsy methods.

Dual Functionality: In addition to taking samples of tissues, EUS-guided biopsy allows doctors to examine tissues and organs closely. This helps in catching problems early and planning treatment more effectively.

Transrectal or Transvaginal Biopsy:

These highly specialized techniques employ ultrasound guidance to obtain biopsy specimens from the prostate or pelvic area (Mascilini, F. *et al.* 2023). Transrectal and transvaginal biopsies are low-risk techniques applied to retrieve tissue samples for diagnostic reasons, especially in the event of abnormality in the prostate or pelvic organs. They are guided by imaging devices such as ultrasound for accuracy and safety.

Transrectal Biopsy: Transrectal biopsy is a procedure usually used to diagnose prostate disorders, such as cancer. In a transrectal biopsy, an ultrasound probe is placed in the rectum to see the prostate gland. A thin needle is then passed through the rectal wall to obtain tissue samples from targeted areas of the prostate. The procedure is usually done under local anaesthesia and lasts approximately 15–30 minutes. It is very effective in identifying prostate cancer and other abnormalities. Mild discomfort or minor bleeding after the procedure may be experienced by patients, but recovery is generally rapid.

Transvaginal Biopsy: This procedure is utilized to diagnose conditions of the female reproductive system, like the uterus, ovaries, or pelvic masses (*Respir. Med. Case Rep.*, 2023). A transvaginal ultrasound probe is placed in the vagina to offer real-time images of the area of interest. A biopsy needle removes tissue samples for examination. This procedure is sometimes done on an outpatient basis and is painless, with minimal to no downtime. It is especially helpful for assessing abnormal growths, thickened endometrium, or other suspicious results.

Both tests are useful diagnostic modalities that provide high accuracy with minimal patient discomfort. They are frequently used instead of more invasive surgical techniques because they are efficient and have less risk of complications. Imaging guidance allows the biopsies to be targeted, minimizing sampling errors.

These methods are important for early detection and treatment planning for many conditions, enhancing patient outcomes and quality of care.

Each is selected according to the location, size, and type of the abnormality being studied.

CONCLUSION:

Ultrasound-guided biopsy is a milestone in diagnostic medicine, with unmatched accuracy, safety, and efficiency. It employs real-time imaging to locate and biopsy abnormal tissues from different organs such as the thyroid, liver, pancreas, breast, and lymph nodes. By directly targeting suspicious regions, ultrasound-guided biopsy reduces trauma to surrounding tissues and delivers high-quality samples for analysis, rendering it a precious asset in the diagnosis and treatment of conditions like cancer, infection, and inflammatory disease. One of the most characteristic strengths of ultrasound-guided biopsy is its patient-friendliness. The procedure is minimally invasive, causing minimal pain to patients in comparison to conventional surgical techniques. It is done through real-time imaging, which provides accurate targeting of abnormalities even in deep or inaccessible locations. Furthermore, the absence of exposure to ionizing radiation, as observed in imaging modalities such as X-rays and CT scans, renders it safer for patients, particularly those that need repeated treatments. Its cost-effectiveness is another key advantage, since it obviates the need for elaborate surgical arrangements and prolonged in-hospital stays, allowing it to be offered to more patients. Breakthroughs like endoscopic ultrasound-guided biopsy have extended the use of this procedure to new dimensions, enabling doctors to reach and diagnose lesions in hitherto difficult locations such as the gastrointestinal tract. Endoscopic ultrasound unites the strengths of conventional ultrasound imaging with the power to maneuver along internal tubes, giving complete visualization and pinpoint accuracy to target suspicious lesions. This technology has tremendously boosted the diagnostic power of ultrasound-guided biopsy, positioning it as a mainstay of contemporary medicine. Though it has fewer drawbacks, ultrasound-guided biopsy still has limitations. The procedure is highly operator-dependent because the skill and experience of the performing physician are crucial in ensuring good imaging and sample collection. Sampling from an incorrect location or not getting a proper sample for analysis may happen from time to time. To avoid this, however, with adequate training, proper practice, and strict adherence to guidelines, such risks can be kept under control. Patient care is a critical component of minimizing complications because careful pre-procedure planning and post-procedure monitoring can keep adverse results at a minimum. Ultrasound-guided biopsy is also an important interface between imaging and pathology. By merging diagnostic imaging with tissue analysis, the procedure achieves a complete

picture of the condition at hand, making it possible for better diagnoses and tailored treatment plans. The feature to integrate real-time imaging with direct tissue sampling allows clinicians to make timely and informed decisions, cutting down on delays in diagnosis and enhancing outcomes for patients. Continued research and advances in technology continue to fine-tune and enhance the application of ultrasound-guided biopsy. Advances in imaging technology, including the development of three-dimensional ultrasound and more advanced software algorithms, will further enhance the accuracy and efficiency of this method. Further, incorporation of artificial intelligence into imaging analysis will be potentially beneficial in diminishing operator dependence and decreasing errors. These developments seek to further improve ultrasound-guided biopsy as safer, more efficient, and more versatile, leading to improved disease management and enhanced patient care. Overall, ultrasound-guided biopsy underscores the revolutionary impact of technology on contemporary healthcare. It is a model for a move toward more accurate, patient-focused medical procedures, making diagnostic interventions both effective and less invasive. By offering accurate results with little pain, this technique has emerged as a primary instrument in the diagnosis and treatment of many conditions.

REFERENCES:

1. Davis, M., Sofer, M., Kim, S.S. and Soloway, M.S., 2002. The procedure of transrectal ultrasound guided biopsy of the prostate: a survey of patient preparation and biopsy technique. *The Journal of urology*, 167(2 Part 1), pp.566-570.
2. Choi, A.Y., Kolb, J., Shah, S., Chahine, A., Hashimoto, R., Patel, A., Tsujino, T., Huang, J., Hu, K.Q., Chang, K. and Samarasena, J.B., 2022. Endoscopic ultrasound-guided portal pressure gradient with liver biopsy: 6 years of endo-hepatology in practice. *Journal of gastroenterology and hepatology*, 37(7), pp.1373-1379.
3. Facciorusso, A., Ramai, D., Conti Bellocchi, M.C., Bernardoni, L., Manfrin, E., Muscatiello, N. and Crinò, S.F., 2021. Diagnostic yield of endoscopic ultrasound-guided liver biopsy in comparison to percutaneous liver

biopsy: a two-center experience. *Cancers*, 13(12), p.3062.

4. Fallenberg, E.M., 2022. Ultrasound-Guided Interventions. In *Breast Imaging: Diagnosis and Intervention* (pp. 143-163). Cham: Springer International Publishing.
5. Bass, E.J., Pantovic, A., Connor, M.J., Loeb, S., Rastinehad, A.R., Winkler, M., Gabe, R. and Ahmed, H.U., 2022. Diagnostic accuracy of magnetic resonance imaging targeted biopsy techniques compared to transrectal ultrasound guided biopsy of the prostate: a systematic review and meta-analysis. *Prostate Cancer and Prostatic Diseases*, 25(2), pp.174-179.
6. Lubner, M.G., Mankowski Gettle, L., Kim, D.H., Ziemlewicz, T.J., Dahiya, N. and Pickhardt, P., 2021. Diagnostic and procedural intraoperative ultrasound: technique, tips and tricks for optimizing results. *The British journal of radiology*, 94(1121), p.20201406.
7. Mascilini, F., Quagliozzi, L., Mirandola, M., Moro, F., D'Alessandris, N., De Felice, F., Testa, A.C., Scambia, G. and Fagotti, A., 2023. Transvaginal ultrasound-guided biopsy in patients with suspicious primary advanced tubo-ovarian carcinoma. *International Journal of Gynecological Cancer*, 33(2), pp.236-242.
8. Takano, Y., Noda, J., Yamawaki, M., Azami, T., Kobayashi, T., Niiya, F., Maruoka, N., Norose, T., Ohike, N., Wakabayashi, T. and Matsuo, K., 2021. Comparative study of an ultrasound-guided percutaneous biopsy and endoscopic ultrasound-guided fine-needle aspiration for liver tumors. *Internal Medicine*, 60(11), pp.1657-1664.

9. Facciorusso, A., Crinò, S.F., Gkolfakis, P., Ramai, D., Lisotti, A., Papanikolaou, I.S., Mangiavillano, B., Tarantino, I., Anderloni, A., Fabbri, C. and Triantafyllou, K., 2022. Endoscopic ultrasound fine-needle biopsy vs fine-needle aspiration for lymph nodes tissue acquisition: a systematic review and meta-analysis. *Gastroenterology Report*, 10, p.goac 062.
10. Madhok, I.K., Parsa, N. and Nieto, J.M., 2022. Endoscopic ultrasound-guided liver biopsy. *Clinics in Liver Disease*, 26(1), pp.127-138.
11. Sountoulides, P., Pyrgidis, N., Polyzos, S.A., Mykoniatis, I., Asouhidou, E., Papatsoris, A., Dellis, A., Anastasiadis, A., Lusuardi, L. and Hatzichristou, D., 2021. Micro-ultrasound-guided vs multiparametric magnetic resonance imaging-targeted biopsy in the detection of prostate cancer: a systematic review and meta-analysis. *The Journal of urology*, 205(5), pp.1254-1262.
12. Smith, S., 1985. Ultrasound-guided biopsy. *Veterinary Clinics of North America: Small Animal Practice*, 15(6), pp.1249-1262.
13. Siddiqui MM, Rais-Bahrami S, Turkbey B, George AK, Rothwax J, Shakir N, Okoro C, Raskolnikov D, Parnes HL, Linehan WM, Merino MJ, Simon RM, Choyke PL, Wood BJ, Pinto PA. Comparison of MR/ultrasound fusion-guided biopsy with ultrasound-guided biopsy for the diagnosis of prostate cancer. *JAMA*. 2015 Jan 27;313(4):390-7. doi: 10.1001/jama.2014.17942. PMID: 25626035; PMCID: PMC4572575.
14. Haider MA, van der Kwast TH, Tanguay J, et al. Combined T2-weighted and diffusion-weighted MRI for localization of prostate cancer. *AJR Am J Roentgenol*. 2007;189(2):323–328. doi: 10.2214/AJR.07.2211.
15. Kaplan I, Oldenburg NE, Meskeel P, Blake M, Church P, Holupka EJ. Real time MRI-ultrasound image guided stereotactic prostate biopsy. *Magn Reson Imaging*. 2002;20(3):295–299. doi: 10.1016/s0730-725x(02)00490-3.
16. Maskey, A. and Bhoomi, K.K., 2014. USG assisted and USG guided percutaneous renal biopsy at Nepal Medical College Teaching Hospital: A three and half years study. *Nepal Med Coll J*, 16(1), pp.26-9.
17. Govindarajan, P. and Keshava, S.N., 2010. Ultrasound-guided omental biopsy: Review of 173 patients. *Indian Journal of Radiology and Imaging*, 20(04), pp.307-309.
18. Li, X. and Kong, L., 2023. Ultrasound versus computed tomography-guided transthoracic biopsy for pleural and peripheral lung lesions: a systematic review and meta-analysis. *Acta Radiologica*, 64(12), pp.2999-3008.
19. Bharti, S.J. and Hoda, W., 2024. Mass for USG-/CT-Guided Biopsy. *Problem Based Learning Discussions in Onco-Anesthesia and Onco-Critical Care*, p.233.
20. Slobodnikova, J., 2022. The benefit of usg in diagnosing radial scars in the context of differential diagnosis of breast lesions. *Ultrasound in Medicine & Biology*, 48, p.S19.
21. Serinsöz, S. and Akturk, R., 2022. Comparison of diagnostic accuracies of USG, MG and MRI modalities defined with BI-RADS Classification System. *Current Medical Imaging Reviews*, 18(9), pp.986-995.
22. Keskin, L., Karahan, D. and Yaprak, B., 2023. Comparison of thyroid fine needle

aspiration biopsy and ultrasonography results. *Medicine*, 102(26), p.e33822.

23. SHOWKAT, M.S., BHOWMIK, B., SHIRIN, M., GHAFOR, N., MAHDI, R., SHARMIN, F. and RAHMAN, M.T., 2021. Approach of USG guided core biopsy in suspicious breast lesions (BI-RADS 4) depicted by USG & mammography in relation to histopathological outcome: a cross sectional study of 100 cases in a tertiary care center. *Bangladesh Journal of Radiology and Imaging Vol*, 29, p.1.

24. Bharti, S.J. and Hoda, W., 2024. Paediatric Case of Mediastinal Mass for USG-/CT-Guided Biopsy. In *Problem Based Learning Discussions in Onco-Anesthesia and Onco-Critical Care* (pp. 233-240). Singapore: Springer Nature Singapore.

25. Baijal, N. and Chandrashekhara, S.H., 2025. Image-Guided Biopsy. In *Textbook of Interventional Radiology* (pp. 415-426). Singapore: Springer Nature Singapore.

26. Demirelli, B., Boynueğri, B., Boztepe, B., Şenol, E.G., Ramazan, A., Canbakan, M., Gümrükçü, G. and Öğütmen, M.B., Frequency and Risk Factors of Ultrasound-Guided Kidney Biopsy.

27. Nguyen, H., Nguyen, X., Pham, L., Nguyen, T., Vo, T., Le, T., Le, B., Nguyen, T. and Le, C., 2024, August. Ultrasound-guided fine-needle aspiration biopsy of thyroid nodules smaller than 10 mm in the maximum diameter: the efficacy and its correlation with tirads classification. In *Endocrine Abstracts* (Vol. 101). Bioscientifica.

28. Sogukpinar, O., Akturk, U.A., Akbay, M.O., Tatlidil, E. and Ernam, D., 2024. Diagnostic value of transthoracic needle biopsy in

lung tumors. *Revista da Associação Médica Brasileira*, 70(3), p.e20231082.

29. Cay, M., Turan, I., Mengen, E., Çimen, A.M., Celebi, Ş.T. and Yüksel, B., 2025. Optimal timing of repeat thyroid fine-needle aspiration biopsy. *Journal of Pediatric Endocrinology and Metabolism*, (0).

30. Dhamija, E., Popat, P.B. and Syed, A., 2025. Breast Interventions: Biopsy and Localization Techniques. In *Textbook of Interventional Radiology* (pp. 515-524). Singapore: Springer Nature Singapore.